

·Original Article·

Association of *USP26* haplotypes in men in Taiwan, China with severe spermatogenic defect

I-Wen Lee¹, Long-Ching Kuan², Chien-Hung Lin³, Hsien-An Pan¹, Chao-Chin Hsu⁴, Yung-Chieh Tsai⁵, Pao-Lin Kuo¹, Yen-Ni Teng⁶

¹Department of Obstetrics & Gynecology, National Cheng Kung University, College of Medicine, Tainan 701, Taiwan, China

²Department of Obstetrics and Gynecology, Kuo General Hospital, Tainan 700, Taiwan, China

³Institute of Basic Sciences, National Cheng Kung University, College of Medicine, Tainan 701, Taiwan, China

⁴Department of Obstetrics and Gynecology, China Medical University, Taichung 404, Taiwan, China

⁵Department of Obstetrics and Gynecology, Chi-Mei Medical Center, Tainan 710, Taiwan, China

⁶Department of Biotechnology, Chia Nan University of Pharmacy and Science, Tainan 717, Taiwan, China

Abstract

Aim: To complete comprehensive haplotype analysis of *USP26* for both fertile and infertile men. **Methods:** Two hundred infertile men with severe oligospermia or non-obstructive azoospermia were subjected to sequence analysis for the entire coding sequences of the *USP26* gene. Two hundred men with proven fertility were genotyped by primer extension methods. Allele/genotype frequencies, linkage disequilibrium (LD) characteristics and haplotypes of fertile men were compared with infertile men. **Results:** The allele frequencies of five single nucleotide polymorphisms (370-371insACA, 494T>C, 576G>A, ss6202791C>T, 1737G>A) were significantly higher in infertile patients than control subjects. The major haplotypes in infertile men were TACCGA (28% of the population), TGCCGA (15%), TACCAA (8%), TGCCAA (6%), TATCAA (5%) and CATCAA (5%). The major haplotypes for the control subjects were TACCGA (58% of the population), CACCGA (7%), CATCGA (6%) and TGCCGA (5%). Haplotypes TGCCGA, TATCAA, CATCAA, CATCGC, TACCAA and TGCCAA were over-transmitted in patients with spermatogenic defect, whereas haplotypes TACCGA, CACCGA, and CATCGA were under-transmitted in these patients. **Conclusion:** Some *USP26* alleles and haplotypes are associated with spermatogenic defect in the Han nationality in Taiwan, China. (*Asian J Androl* 2008 Nov; 10: 896–904)

Keywords: single nucleotide polymorphism; *USP26* gene; spermatogenic defect; linkage disequilibrium

1 Introduction

Infertility affects one in 10 couples worldwide. In roughly half of these couples, the male factor is implicated. Although there are many factors that contribute to male infertility, the cause cannot be identified for the majority of infertile men [1]. In recent years, there

Correspondence to: Prof. Yen-Ni Teng, Department of Biotechnology, Chia Nan University of Pharmacy and Science, 60 Erh-Jen Road, Sec. 1, Pao-An, Jen-Te Hsiang, Tainan 717, Taiwan, China.
Tel: +886-6-266-4911 ext. 2549 Fax: +886-6-266-2135
E-mail: tengyenni1@yahoo.com.tw
Received 2008-02-20 Accepted 2008-07-03

has been an intensive search for genetic causes of male infertility. Screening with markers on the long arm of the human Y chromosome has detected Yq microdeletions in 5%–15% of males with spermatogenic defect. It is believed that many genes located on the long (q) arm of the Y chromosome are required for spermatogenesis. This region, banding in q11.23 of the Y chromosome, includes the azoospermia factor (AZF) locus, which contains a gene or genes that are required for normal spermatogenesis [2, 3]. Other genetic factors that have been reported to be involved in spermatogenic defect include mutations at the mitochondrial DNA polymerase locus, a polymorphism of the cytochrome P4501A1 gene, mutations of the follicle-stimulating hormone receptor gene (*FSH*), the deleted in azoospermia-like gene (*DAZL*), the synaptonemal complex protein 3 gene (*SYCP3*), human *BOULE*, casein kinase 2 alpha genes, dystrophin myotonia protein kinase gene (*DMPK*), DNA polymerase gamma (*POLG*), methylenetetrahydrofolate reductase (*MTHFR*) and follicle-stimulating hormone receptor (*FSHR*) [4–16].

USP26 appears to be an attractive candidate for a sterile gene considering its unique expression pattern. *USP26* comprises a single exon and is located on Xq26.2. Its messenger RNA (mRNA) sequence is 2 794 bp long and its protein consists of 913 amino acids (Genbank: NM_031907.1). Its mouse homolog was found to be only expressed in spermatogonia [17]. *USP26* belongs to a large family of deubiquitinating enzymes (DUB) [18]. Deubiquitination of macromolecules by DUB, including ubiquitin proteases, can rescue macromolecules from degradation through substrate-specific, N-terminal-dependent, enzymatic reaction [19, 20]. Previous studies have addressed the importance of the ubiquitin pathway during mammalian fertilization, including acrosomal function and spermatozoa–zona pellucida (ZP) penetration [21,22]. During spermatogenesis, deubiquitination enzymes have been shown to be involved in the regulation of protein turnover (e.g., replacement of histones by protamine, germ cell apoptosis, mitotic proliferation and differentiation of spermatogonial stem cells) [23–28]. Several papers have been published on the role of *USP26* in male infertility. Stouffs *et al.* [29] analyzed patients with various histological patterns of spermatogenic defect (Sertoli cell-only syndrome and maturation arrest) for the presence of mutations in *USP26*. They found some Sertoli cell-only syndrome (SCOS) patients had the genotype of 370-371insACA (rs3840975), 494T>C and 1423C>T (rs41299088) [29]. A study by Paduch

also provided evidence linking *USP26* mutation to male infertility [30]. Although these two studies suggested that a specific genetic cluster (370-371insACA, 494T>C and 1423C>T) might be associated with testicular dysfunction, two other studies showed that this genetic cluster was not restricted to men with testicular dysfunction [31, 32]. In a recent publication on *USP26*, the authors identified several novel mutations in the *USP26* gene that might cause spermatogenesis impairment in a small group of infertile men ($n = 41$) [33]. However, meta-analysis of previous case-control studies (in total, 544 patients and 1 705 controls) revealed no significant association of the 370-371insACA, 494T>C and 1423C>T genotype with male infertility [14, 29–32]. Further studies based on a large group of patients with diversified ethnic backgrounds will be valuable to bring more insight into the role of *USP26* in male infertility.

In the present study, we set out to analyze genetic variants and haplotypes of *USP26* for both fertile and infertile men by sequence analysis for the entire coding region of *USP26*. We found some *USP26* genetic variants and haplotypes were overt-transmitted and some were under-transmitted in patients with severe spermatogenic defect in the Han nationality in Taiwan, China. We also identified some novel genetic variants or mutations in patients. Our result supports important roles of *USP26* in human spermatogenesis.

2 Materials and methods

2.1 Subjects

The study was approved by the Institutional Review Board of National Cheng Kung University Medical Centre (Taiwan, China). From January 2001 to December 2005, 200 infertile men presenting with severe oligozoospermia (spermatozoa count $< 5 \times 10^6/\text{mL}$) or non-obstructive azoospermia and 200 men with proven fertility (control group) were enrolled. All of the control subjects had fathered at least two children within 5 years without assisted reproductive technologies. The paternal relationship between control subjects and their offspring was confirmed using the AmpF/STR-Profiler-Plus zygosity determination system (Applied Biosystems, Foster City, CA, USA). All patients and control subjects belonged to the Han nationality in Taiwan, the major ethnic group in China (making up more than 95% of this province's population). The patients were evaluated according to the protocol described in our previous studies [6, 34]. In

brief, all patients underwent comprehensive characterization, including a detailed history, physical examination, at least two semen analyses, hormone assays (luteinizing hormone [LH], follicle stimulating hormone [FSH], prolactin [PRL] and testosterone [T]), karyotyping and a molecular test for Y-chromosomal deletions. Patients suspected to have non-obstructive azoospermia were advised to undergo bilateral testicular biopsies. Serum levels of FSH, LH, PRL and T were measured using commercial radioimmunoassay kits: Coat-A-Count FSH IRMA, Coat-A-Count LH IRMA, Coat-A-Count PRL IRMA and IMMULITE Total Testosterone (Diagnostic Products, Los Angeles, CA, USA). Chromosome analysis was performed using the GTG method (G-banding by Trypsin-Giemsa technique). Molecular analysis of Y-chromosomal deletions included a combination of 16 gene-based primers, as described previously [34]. Patients with any identifiable cause of male infertility, including congenital bilateral absence of vas deference (CBAVD), cryptorchidism, varicocele, diabetes mellitus or hypertension, or with history that may affect spermatogenesis (e.g., orchitis, trauma, malignancies, *etc.*) were excluded from the study group. No significant difference was found between age, serum LH, estradiol and T concentrations of infertile men and control subjects. Sperm count and FSH concentrations were significantly higher in patients compared with the control group ($P < 0.005$). Of 200 study subjects, 137 had severe oligospermia and 63 had azoospermia.

2.2 Sequence analysis of patients

Genomic DNA samples of patients were subjected to sequence analysis for the entire coding sequences of

the *USP26* gene. Genomic DNA was extracted from peripheral blood samples using a Puregene DNA isolation kit (Gentra, Minneapolis, MN, USA). *USP26* genomic sequence (AF285593) was obtained from the NCBI web site (<http://www.ncbi.nlm.nih.gov>). The entire gene (2 794 bp) was divided into five overlapping fragments ranging from 409 to 600 bp in length. To amplify partial fragments of the *USP26* exon, polymerase chain reaction (PCR) reactions were performed in 20 μ L volumes containing 200 ng of genomic DNA, 10 mmol/L Tris-HCl (pH 8.3), 50 mmol/L KCl, 0.1% Triton X-100, 200 μ mol/L deoxyribonucleotide triphosphates (dNTP), 100 pmol of each primer and 1 U *Taq* DNA polymerase (Promega Corp., Madison, WI, USA). The primer sets for *USP26* are described in (Table 1). PCR amplification was performed in an automated thermal cycler (OmniGene Thermal Cycler; Hybaid Ltd., Ashford Middlesex, UK). PCR products were sequenced to identify mutations or polymorphisms. Sequence analysis was performed with an automatic sequencer (ABI 377, Applied Biosystems/PE).

2.3 Genotyping for fertile controls

Genomic DNA samples of the control subjects were subjected to genotyping by primer extension methods. The amplicons were amplified in a multiplex fashion and each 20 μ L reaction consisted of 50 ng of genomic DNA, 10 pmol of primers, 4.0 mmol/L $MgCl_2$, 0.2 mmol/L dNTP and 0.5 U of AmpliTaq Gold DNA polymerase (Applied Biosystems). The cycling conditions were: 95°C for 5 min, followed by 35 cycles at 95°C for 1 min, 55°C for 1 min and 72°C for 1 min, with a final extension for

Table 1. Primer sets used for PCR and DNA sequencing of the *USP26* gene. ^aThe nucleotide sequences is according to AF285593. ^bThe lengths of polymerase chain reaction product size.

Fragment	Primer pairs	Primer sequence ^a	Product size (bp) ^b
1	Usp26-1F	CTTCAAGATTATCAATAATCGG	680
	Usp26-1R	AATCTGCCTTGGATTCTTG	
2	Usp26-2F	GCTCTCATCTAGCTCAGAGA	687
	Usp26-2R	CCAAATTCACCTTTTAGGCTTCC	
3	Usp26-3F	GCTCATGAGTTTTAGCTCACT	682
	Usp26-3R	GTTCAGACTCCTTATCTGATC	
4	Usp26-4F	CATCAGTGTATCATGGCCTG	562
	Usp26-4R	TTGTGTGCCCTGGGTGCC	
5	Usp26-5F	CGGTATGAGAATCTGTGAAC	511
	Usp26-5R	TTCCTTCTGAAGGGTCTCCTC	

10 min at 72°C. Each allele was measured by primer extension and SNaPshot chemistry (Applied Biosystems). Multiplex PCR products (3 µL) were treated with 2.5 U of shrimp alkaline phosphatase (SAP; Amersham Pharmacia Biotech, Uppsala, Sweden) and 2 U of exonuclease I (BM Biochemica, Mannheim, Germany) in a 10 mL reaction volume for 1 h at 37°C. SNaPshot multiplex PCR was performed in a 10 µL reaction volume containing 1.25 µL of SNaPshot Multiplex Ready Reaction Mix (Applied Biosystems), 4 µL of SAP/exonuclease I-treated PCR products, and 1 to 3 pmol of each SNaPshot primer. Thermal cycling for SNaPshot reactions consisted of 25 cycles at 96°C for 10 s, 50°C for 5 s and 60°C for 30 s. The SNaPshot multiplex reaction was then treated by SAP in an 11 µL reaction volume containing 10 µL of SNaPshot multiplex PCR product and 1 U of SAP. The samples were analyzed using an ABI 3100 genetic analyzer (Applied Biosystems) and the allele determination was carried out with the Genotyper 3.7 program (Applied Biosystems).

2.4 Statistical analysis

Tests for association with single markers and haplotypes in control samples were performed using the χ^2 test. $P < 0.05$ was considered statistically significant. The relative risk of spermatogenic defect was estimated from logistic odds ratios (OR) and 95% confidence intervals (CI) in multivariate analysis. Tests for haplotype association with spermatogenic defect were performed using EHPLUS software [35], and statistical significance was estimated using the permutation test PMPLUS [36]. Associations between haplotypes and male infertility were analyzed using linear regression models. The linkage disequilibrium (LD) coefficient (D') between each pair of single nucleotide polymorphisms (SNPs) was calculated using the ldmax program within the GOLD software package [37].

3 Results

3.1 Polymorphisms in men in Taiwan, China

The coding sequences of the *USP26* gene were examined by sequencing analysis for the presence of mutations of sequence variants in infertile men. The sequence from GenBank (AF285593) was used as the reference sequence and the A of ATG of the start codon was numbered + 1. In total, seven genetic variants were observed, including one insertion variant 370–371insACA

(rs3840975), causing a threonine insertion in amino acid position 121, and six SNPs, 494T>C, 576G>A (rs41304540), ss6202791C>T, 1423C>T (rs41299088), 1737G>A and 2202A>C. All of these genetic variants were available in the web site (<http://www.ncbi.nlm.nih.gov/>) and peer-review articles [29–32]. One substitution, 576G>A (E192), does not alter the amino acid sequence: a glutamic acid at position 192. The other five SNP were predicted to create amino acid alteration: 494T>C changes a leucine into a serine (L165S); ss6202791C>T changes a serine into a phenylalanine (S600F); 1423C>T substitutes a histidine for a tyrosine (H475Y); 1737G>A substitutes a methionine for a isoleucine (M579I); and 2202A>C substitutes a lysine for an asparagine (K734N). The following substitutions occurred only once in the patients group but not in the fertile controls group: 393A>T (S131S); 468T>G (Q156H); 1976C>T (T659M); 2144A>T (N715I); 2182A>T (I128F); 2195T>C (F732S); 2204T>G (V735G); 2239A>T (I747F); 2247A>C (E749D); 2250A>G (Q750Q); and 2271T>C (P757P). Except for 1976C>T (T659M), these substitutions have not been reported [30]. The substitution of 520T>G and 565T>G is predicted to cause premature stop codon at codon 175 and 189, respectively. Haplotype analyses ruled out that patients carrying these variant sites occurred only once in the infertility group.

3.2 Allelic and genotypic frequencies in fertile and infertile men

Allelic frequencies and odds ratios for each sequence variant of *USP26* are shown in Table 2. Of the seven variants, 370–371insACA, 494T>C, 576G>A, ss6202791C>T and 1737G>A are significantly associated with spermatogenic defect. The P -value for these five variants was less than 0.05 and the odds ratios were larger than 1. For example, the P -values for 370–371insACA and 576G>A are less than 0.0001 and the OR are 35.86 and 4.095 (Table 2).

3.3 Reproductive effect of the 70–371insACA, 494T>C, 1423C>T genetic cluster and its derivation

The ancestral *USP26* cluster [370–371insACA, 494T>C, 1423C>T] and its first derivation only occur in the group of infertile men. In contrast, the second derivation only occurs in the group of fertile men (Table 3). The frequency of the [370–371insACA, 494T>C, 1423C>T] cluster is 3% in the infertile group but none in fertile men.

Table 2. Allelic frequencies of seven *USP26* sequence variants in fertile and infertile men. Allele frequencies were compared between infertile men and control subjects. SNPs, single nucleotide polymorphisms; OR, odds ratio; CI, confidence interval.

SNPs	Allele	Allele frequency		P-value (χ^2)	OR	95% CI
		Control (n = 200) n (%)	Spermatogenic defect (n = 200) n (%)			
370-371insACA	No ins	200 (100)	184 (92)	< 0.0001	35.860	2.135–602.400
	insACA	0 (0)	16 (8)			
494T>C	T	164 (82)	146 (73)	0.0311	1.685	1.046–2.715
	C	36 (18)	54 (27)			
576G>A	A	172 (86)	120 (60)	< 0.0001	4.095	2.510–6.682
	G	28 (14)	80 (40)			
ss6202791C>T	C	162 (81)	144 (72)	0.0338	1.658	1.037–2.651
	T	38 (19)	56 (28)			
1423C>T	C	192 (96)	186 (93)	0.1882	1.806	0.740–4.408
	T	8 (4)	14 (7)			
1737G>A	G	172 (86)	120 (60)	< 0.0001	4.095	2.510–6.682
	A	28 (14)	80 (40)			
2202A>C	A	196 (98)	194 (97)	0.5218	1.515	0.421–5.456
	C	4 (2)	6 (3)			

Table 3. The genotype frequency of the 370-371insACA, 494T>C and 1423C>T cluster, its first and second derivative in fertile and infertile men. ^aAnalyzed for statistical significance by Fisher's exact test. ^bAccording to [37]. OR, odds ratio.

	370-371ins ACA	494T>C	1423C>T	Spermatogenic defect n (%)	Controls n (%)	P-value ^a	OR
Ancestral ^b	Yes	C	C	8 (4%)	0 (0%)	0.0073	17.71
1st derivation ^b	Yes	C	T	6 (3%)	0 (0%)	0.0301	13.40
2nd derivation ^b	No	T	T	0 (0%)	4 (2%)	0.1231	0.1089

3.4 Linkage disequilibrium (LD) between different SNPs

(Table 4) shows the pairwise LD (D') between different alleles. For patients with spermatogenic defect, 2202A>C was found to be in complete LD with 370-371insACA, 494T>C, 576G>A and 1423C>T ($D' = 1$). 494T>C had strong LD with 370-371insACA and 1423C>T ($D' = 0.829$ and 0.804 respectively). For control subjects, 370-371insACA was found to be in weak LD with the other six variants ($D' = 0$). Except 2202A>C, no strong linkage disequilibrium ($D' > 0.8$) was observed for these seven variants in the control group. Therefore, the LD characteristic of infertile men was different from that of control subjects.

3.5 Haplotype analysis

The major haplotypes (defined as haplotype frequency

in either group $\geq 5\%$) were TACCGA (28% of the population), TGCCGA (15%), TACCAA (8%), TGCCAA (6%), TATCAA (5%) and CATCAA (5%) for patients. The capital letters represented 494T>C, 576G>A, ss6202791C>T, 1423C>T, 1737G>A and 2202A>C from left to right, respectively. The major haplotypes for the control subjects were TACCGA (58% of the population), CACCGA (7%), CATCGA (6%) and TGCCGA (5%). Six of the nine major haplotypes showed significant differences in frequency between infertile men and control subjects (Table 5). Haplotypes TGCCGA, TATCAA, CATCAA, CATCGC, TACCAA and TGCCAA were over-transmitted in patients with severe spermatogenic defect, whereas haplotype TACCGA, CACCGA and CATCGA were under-transmitted in these patients. Of 134 patients with major haplotypes, only nine azoospermic men

Table 4. Pairwise linkage disequilibrium (D') in fertile and infertile men. Estimates in the upper right are for infertile men and in the lower left are for control subjects.

	370-371insACA	494T>C	576G>A	ss6202791C>T	1423C>T	1737G>A
370-371insACA	—	0.829	0.375	0.107	0.534	0.583
494T>C	0	—	0.136	0.331	0.804	0.259
576G>A	0	0.206	—	0.107	0.762	0.042
ss6202791C>T	0	0.451	0.206	—	0.008	0.464
1423C>T	0	0.390	0.128	0.691	—	0.286
1737G>A	0	0.206	0.336	0.206	0.128	—
2202A>C	0	1	1	0.383	1	1

Table 5. Major haplotypes of fertile and infertile men. *Analyzed for statistical significance by Fisher's exact test and $P < 0.05$ was considered statistically significant. OR, odds ratio; CI, confidence interval.

Patterns	Haplotype							Spermatogenic defect vs. controls				
	370-371 insACA	494T >C	576G >A	ss6202791C >T	1423C >T	1737G >A	2202A >C	Spermatogenic defect	Controls	P-value ^a	OR	95% CI
1	No	T	A	C	C	G	A	56 (28%)	116 (58%)	< 0.0001	0.282	0.186–0.428
2	No	T	G	C	C	G	A	30 (15%)	10 (5%)	0.0013	3.353	1.591–7.064
3	No	C	A	C	C	G	A	2 (1%)	14 (7%)	0.0035	0.134	0.030–0.599
4	No	T	A	T	C	A	A	10 (5%)	2 (1%)	0.0358	5.211	1.127–24.100
5	No	C	A	T	C	A	A	10 (5%)	2 (1%)	0.0358	5.211	1.127–24.100
6	No	C	A	T	C	G	C	8 (4%)	0 (0%)	0.0073	17.710	1.014–309.100
7	No	T	A	C	C	A	A	16 (8%)	8 (4%)	0.1391	2.087	0.872–4.995
8	No	T	G	C	C	A	A	12 (6%)	8 (4%)	0.4923	1.532	0.612–3.833
9	No	C	A	T	C	G	A	8 (4%)	12 (6%)	0.4923	0.653	0.261–1.633

Table 6. Clinical features of patients with major haplotypes. HS, Hypospermatogenesis; MA, maturation arrest; SCOS, Sertoli cell-only syndrome.

Haplotype	Number of patients	Severe oligospermia (No.)	Azoospermia (No.)	Testicular histology (Biopsy: No.)
1	56	48	8	HS: 1, MA: 1
2	30	26	4	MA: 2
4	10	4	6	SCOS: 1
5	10	6	4	MA: 1
7	16	8	8	HS: 1, SCOS: 1
8	12	10	2	HS: 1
Total number of patients	134	102	32	9

had testicular histology available. We found their testicular histological features were highly variable, ranging from hypospermatogenesis, MA (maturation arrest at spermatocyte or spermatid) to SCOS (Table 6).

4 Discussion

In the present study, we found allele frequencies of five SNPs (370-371insACA, 494T>C, 576G>A,

ss6202791C>T and 1737G>A) were significantly higher in infertile patients than those in control subjects (Table 2). As compared with previous studies, our data represent a comprehensive coverage of USP26 genotypes by sequence analysis for the entire coding region. By sequence analysis for the entire coding region, we also discovered some novel variants (e.g., 393A>T, 468T>G, 520T>G, 565T>G, 2144A>T, 2182A>T, 2195T>C, 2204T>G, 2239A>T, 2247A>C, 2250A>G and 2271T>C). These substitutions occurred only once in the patients group but not in the fertile men. Genotyping for more subjects will be required to determine if these substitutions are rare SNP or true mutation.

Although our study is in accordance with two previous studies that described association of genetic cluster (370-371insACA, 494T>C, 1423C>T) with male infertility [29, 30, 32], the role of 370-371insACA, 494T>C and 1423C>T genotypes remains uncertain. Meta-analysis of four studies (a total of 544 patients and 1705 controls) revealed no significant association of the 370-371insACA, 494T>C and 1423C>T genotypes with male infertility (14, 29–32). After incorporating our data (a total of five studies with 744 patients and 1905 controls), meta-analysis still revealed negative association (OR: 0.778; 95% CI: 0.484–1.252). We speculate that the uncertainty could be accounted for by genetic backgrounds of the enrollees in different studies. The frequencies of the 370-371insACA, 494T>C and 1423C>T genotypes seem highly varied demographically even in the Chinese population. We could hardly detect the 370-371insACA, 494T>C and 1423C>T genotypes in fertile men of the Han nationality in Taiwan, China (Table 3). The frequency of ancestral cluster in our study is 4% in infertile men and 0% in fertile controls. The overall frequency in fertile men is 2% for the Han nationality in Taiwan, China, much lower than that reported in the Chinese Han (8%) [31]. The genes involved in reproduction have been shown to be under strong positive selection [38]. The Han immigrated from Mainland China to Taiwan Province relatively recently – within the past several hundred years [39]. The low prevalence of 370-371insACA, 494T>C and 1423C>T genotypes in the Han nationality in Taiwan, China could be interpreted as a result of migration followed by genetic drift.

Recently, LD has been used to track down candidate genes in association studies, in which the disease variants are detected through the presence of linkage signal at

nearby sites [40]. Patterns of LD are also important for unraveling the evolutionary history of humans, including identification of demographic effect and the detection of natural selection [41]. We found different LD characteristics of USP26 for fertile and infertile men. This finding may imply distinct genetic backgrounds for fertile and infertile men. In infertile men, 2202A>C was found to be in complete LD with 370-371insACA, 494T>C, 576G>A and 1423C>T ($D' = 1$). The 494T>C allele had strong LD with 370-371insACA and 1423C>T ($D' = 0.829$ and 0.804 , respectively) (Table 4). The major haplotypes for patients with spermatogenic defect were TACCGA (28% of the population), TGCCGA (15%), TACCAA (8%), TGCCAA (6%), TATCAA (5%) and CATCAA (5%). These six haplotypes accounted for 67% of the patient population. The major haplotypes for control subjects were TACCGA (58% of the population), CACCGA (7%), CATCGA (6%) and TGCCGA (5%), which altogether constitute 76% of the control population. It appeared that haplotypes TGCCGA, TATCAA, CATCAA, CATCGC, TACCAA and TGCCAA conferred susceptibility to spermatogenic defect, whereas haplotype TACCGA, CACCGA and CATCGA were protective against spermatogenic defect (Table 5). It is noteworthy that, of the nine deleterious haplotypes, five belonged to minor haplotype (< 5%) of the fertile men, reflecting negative selection in the population. It is interesting to note occurrence of various testicular phenotypes in men with deleterious USP26 haplotypes. This observation is in accordance with a complex disease model for spermatogenic defect [42]. Male infertility represents a classical example of a complex disease with substantial genetic contribution. USP26 genotype is among one of the many factors that may contribute to spermatogenic defect in humans.

Taken together, we showed association of specific genetic variants with severe spermatogenic defect in the Taiwanese Han. We identified haplotypes of USP26 and found association of specific haplotypes with impaired sperm production for the first time. Our result seems to support important roles of USP26 in human spermatogenesis. Considering contradictory data about the clinical implication of specific USP26 genotypes, USP26 may be just among one of the many factors contributing to spermatogenic defect. Other factors that are involved in diverse regulatory pathways of human spermatogenesis may jointly modify the effect of USP26.

Acknowledgment

The expenses for the publication of this paper were supported by grants from the National Science Council of Taiwan, China (No. NSC- 91-2314-B-006-149, NSC 91-3112-B-006-008, NSC 92-3112-B-006-002, NSC 93-3112-B-006-004, NSC 93-2314-B-006-078, NSC94-2320-B-041-002 and NSC95-2320-B-041-006).

References

- Huynh T, Mollard R, Trounson A. Selected genetic factors associated with male infertility. *Hum Reprod Update* 2002; 8: 183–98.
- Reijo R, Alagappan RK, Patrizio P, Page DC. Severe oligospermia resulting from deletions of spermatogenic defect factor gene on Y chromosome. *Lancet* 1996; 347: 1290–3.
- Pryor JL, Kent-First M, Muallem A, van Bergen AH, Nolten WE, Meisner L, *et al.* Microdeletions in the Y chromosome of infertile men. *N Engl J Med* 1997; 336: 534–9.
- Trannainen JS, Aittomäki K, Min J, Vaskivuo T, Huhtaniemi IT. Men homozygous for an inactivating mutation of the follicle-stimulating hormone (FSH) receptor gene present variable suppression of spermatogenesis and infertility. *Nat Genet* 1997; 15: 205–6.
- Fritsche E, Schuppe HC, Dohr O, Ruzicka T, Gleichmann E, Abel J. Increased frequencies of cytochrome P4501A1 polymorphisms in infertile men. *Andrologia* 1998; 30: 125–8.
- Teng YN, Lin YM, Lin YH, Tsao SY, Hsu CC, Lin SJ, *et al.* Association of a single nucleotide polymorphism of the deleted-in- azoospermia-like (DAZL) gene with susceptibility to spermatogenic defect. *J Clin Endocrinol & Metab* 2002; 87: 5258–64.
- Miyamoto T, Hasulke S, Yorgev L, Maduro MR, Ishikawa M, Westphal H, *et al.* Azoospermia in patients heterozygous for a mutation in SYCP3. *J Clin Endocrinol & Metab* 2002; 87: 5258–64.
- Truong BN, Moses EK, Armes JE, Venter DJ, Baker HW. Searching for candidate genes for male infertility. *Asian J Androl* 2003; 5(2):137–47.
- Lin YW, Hsu LC, Kuo PL, Huang WJ, Chiang HS, Yeh SD, *et al.* Partial duplication at AZFc on the Y chromosome is a risk factor for impaired spermatogenesis in Han Chinese in Taiwan. *Hum Mutat* 2007; 28: 486–94.
- Ferlin A, Raicu F, Gatta V, Zuccarello D, Palka G, Foresta C. Male infertility: role of genetic background. *Reprod Biomed Online* 2007; 14: 734–45.
- Wu B, Lu NX, Xia YK, Gu AH, Lu CC, Wang W, *et al.* A frequent Y chromosome b2/b3 subdeletion shows strong association with male infertility in Han-Chinese population. *Hum Reprod* 2007; 22: 1107–13.
- Zhang F, Li Z, Wen B, Jiang J, Shao M, Zhao Y, *et al.* A frequent partial AZFc deletion does not render an increased risk of spermatogenic impairment in East Asians. *Ann Hum Genet* 2006; 70: 304–13.
- A ZC, Yang Y, Zhang SZ, Li N, Zhang W. Single nucleotide polymorphism C677T in the methylenetetrahydrofolate reductase gene might be a genetic risk factor for infertility for Chinese men with azoospermia or severe oligozoospermia. *Asian J Androl* 2007; 9: 57–62.
- Tüttelmann F, Rajpert-De Meyts E, Nieschlag E, Simoni M. Gene polymorphisms and male infertility – a meta-analysis and literature review. *Reprod Biomed Online* 2007; 15: 643–58.
- Gu AH, Liang J, Lu NX, Wu B, Xia YK, Lu CC, *et al.* Association of *XRCC1* gene polymorphisms with idiopathic azoospermia in a Chinese population. *Asian J Androl* 2007; 9: 781–6.
- Yang Y, Xiao CY, A ZC, Zhang SZ, Li X, Zhang SX. DAZ1/DAZ2 cluster deletion mediated by gr/gr recombination *per se* may not be sufficient for spermatogenesis impairment: a study of Chinese normozoospermic men. *Asian J Androl* 2006; 8: 183–7.
- Wang PJ, McCarrey JR, Yang F, Page DC. An abundance of X-linked genes expressed in spermatogonia. *Nat Genet* 2001; 27: 422–6.
- Glickman MH, Ciechanover A. The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev* 2002; 82: 373–428.
- Wilkinson KD. Regulation of ubiquitin-dependent processes by deubiquitinating enzymes. *FASEB J* 1997; 11: 1245–56.
- Wing SS. Deubiquitinating enzymes - the importance of driving in reverse along the ubiquitin-proteasome pathway. *Int J Biochem Cell Biol* 2003; 35: 590–605.
- Yi YJ, Manandhar G, Sutovsky M, Li R, Jonáková V, Oko R, *et al.* Ubiquitin C-terminal hydrolase-activity is involved in sperm acrosomal function and anti-polyspermy defense during porcine fertilization. *Biol Reprod* 2007; 77: 780–93.
- Yi YJ, Manandhar G, Oko RJ, Breed WG, Sutovsky P. Mechanism of sperm-zona pellucida penetration during mammalian fertilization: 26S proteasome as a candidate egg coat lysin. *Soc Reprod Fertil Suppl* 2007; 63: 385–408.
- Baarends WM, Roest HP, Grootegoed JA. The ubiquitin system in gametogenesis. *Mol Cell Endocrinol* 1999; 151: 5–16.
- Baarends WM, van der Laan R, Grootegoed JA. Specific aspects of the ubiquitin system in spermatogenesis. *J Endocrinol Invest* 2000; 23: 597–604.
- Bebington C, Doherty FJ, Fleming SD. The possible biological and reproductive functions of ubiquitin. *Hum Reprod Update* 2001; 7: 102–11.
- Wang YL, Liu W, Sun YJ, Kwon J, Setsuie R, Osaka H, *et al.* Overexpression of ubiquitin carboxyl-terminal hydrolase L1 arrests spermatogenesis in transgenic mice. *Mol Reprod Dev* 2006; 73: 40–9.
- Wright A, Reiley WW, Chang M, Jin W, Lee AJ, Zhang M, *et al.* Regulation of early wave of germ cell apoptosis and spermatogenesis by deubiquitinating enzyme CYLD. *Dev Cell* 2007; 13:705–16.
- Baarends WM, Roest HP, Grootegoed JA. The ubiquitin system in gametogenesis. *Mol Cell Endocrinol* 1999; 151: 5–16.

- 29 Stouffs K, Lissens W, Tournaye H, Van Steirteghem A, Liebaers I. Possible role of USP26 in patients with severely impaired spermatogenesis. *Eur J Hum Genet* 2005; 13: 336–40.
- 30 Paduch DA, Mielnik A, Schlegel PN. Novel mutations in testis-specific ubiquitin protease 26 gene may cause male infertility and hypogonadism. *Reprod Biomed Online* 2005; 10: 747–54.
- 31 Ravel C, El Houate B, Chantot S, Lourenço D, Dumaine A, Rouba H, *et al.* Haplotypes, mutations and male fertility: the story of the testis-specific ubiquitin protease USP26. *Mol Hum Reprod* 2006; 12: 643–6.
- 32 Stouffs K, Lissens W, Tournaye H, van Steirteghem A, Liebaers I. Alterations of the *USP26* gene in Caucasian men. *Int J Androl* 2006; 29: 614–7.
- 33 Zhang J, Qiu SD, Li SB, Zhou DX, Tian H, Huo YW, *et al.* Novel mutations in ubiquitin-specific protease 26 gene might cause spermatogenesis impairment and male infertility. *Asian J Androl* 2007; 9: 809–14.
- 34 Lin YM, Lin YH, Teng YN, Hsu CC, Lin JSN, Kuo PL. Gene-based screening for Y-chromosome deletions in Taiwanese men presenting with spermatogenic defect. *Fertil Steril* 2002; 77: 897–903.
- 35 Xie X, Ott J. Testing linkage disequilibrium between a disease gene and marker loci. *Am J Hum Genet* 1993; 53: 1107–9.
- 36 Zhao J, Curtis D, Sham PC. Model-free analysis and permutation tests for allelic associations. *Hum Hered* 2000; 50: 133–9.
- 37 Abecasis GR, Cookson WO. Gold-graphical overview of linkage disequilibrium. *Bioinformatics* 2000; 16: 182–30.
- 38 Vallender EJ, Lahn BT. Positive selection on the human genome. *Hum Mol Genet* 2004; 13 Spec No 2: R245–54.
- 39 Ge JX, Wu SD, Chao SJ. *The Migration History of China*. Fuzhou: Fujian People Press; 1997 (In Chinese).
- 40 Wall JD, Pritchard JK. Haplotype blocks and linkage disequilibrium in the human genome. *Nat Rev Genet* 2003; 4: 587–97.
- 41 Coop G, Przeworski M. An evolutionary view of human recombination. *Nat Rev Genet* 2007; 8: 23–34.
- 42 Ferlin A. Male infertility: role of genetic background. *Reprod BioMed Online* 2007; 14: 734–45.

Edited by Gordon Baker